

Women's Sexual Health Survey

Over the last 4 weeks, how would you rate the following:

1. When you have sexual stimulation or intercourse how often do you reach a clitoral orgasm?
0 = rarely // 1 = sometimes // 2 = often // 3 = always
2. When you have sexual stimulation or intercourse how difficult is it to reach a clitoral orgasm?
0 = extremely difficult // 1 = moderately difficult // 2 = Not difficult
3. When you have sexual stimulation or intercourse, how many clitoral orgasms do you typically achieve?
0 = none // 1 = one // 2 = two plus
4. When you reach clitoral orgasm or climax, how satisfied are you with the overall intensity?
0 = never happens // 1 = mildly satisfied // 2 = satisfied
5. How would you rate your level of sexual arousal during intercourse or sexual activity?
0 = very little arousal // 1 = moderate arousal // 2 = extremely aroused

Scoring Instructions:

Add the numbers up to determine whether the patient characterizes symptoms of clitoral sexual dysfunction and whether or not Cliovana is recommended.

13-15	//	No clitoral sexual dysfunction – Treatment not necessary
10-13	//	Mild clitoral sexual dysfunction – Treatment recommended
6-9	//	Mild to Moderate clitoral sexual dysfunction – Treatment recommended
3-5	//	Moderate clitoral sexual dysfunction – Treatment recommended
0-2	//	Severe clitoral sexual dysfunction – Treatment recommended

The purpose of the survey is to determine whether or not a patient is a candidate for the Cliovana treatment and to also serve as a baseline before treatments begin.

- The survey should be conducted **THREE TIMES**: (1) once during the consultation, (2) after the series 4 Cliovana treatments, and (3) 12 weeks post the patients 4th Cliovana treatment.